

UNCLASSIFIED

AD NUMBER
AD845941
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; JAN 1969. Other requests shall be referred to Commanding Officer, Fort Detrick, Attn: SMUFD-AE-T, Frederick, MD 21701.
AUTHORITY
SMUFD, D/A, 17 Feb 1972

THIS PAGE IS UNCLASSIFIED


AD845941

TRANSLATION NO. 2370

DATE: Jan 6, 1969

DDC AVAILABILITY NOTICE

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Commanding Officer, Fort Detrick, ATTN: SMUPD-AE-T, Frederick, Md. 21701



DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

**Best
Available
Copy**

URINARY HYDROXYPROLINE AND CIRCULATING HGH PRELIMINARY
FINDINGS IN SOME ENDOCRINE DISEASES

Folia Endocrinologica (Endocrinological
Folios), Vol 20, No 4, July-August 1967,
pp 420-441

O. Recchia, L. Sereno, and
I. Isidori

During the approximately 50 years since the first research on urinary polypeptides (77), at least 47 different peptidic bodies have been identified in a rather voluminous and recent series of studies (8, 12, 16, 17, 73, 152, 178).

The volume of the excretion and the qualitative differences recorded under various physical and pathological conditions today seem to confer upon these nitrogenous products a rather significant value as an expression of the protein metabolism. During these past years, special interest was devoted to the study of the urinary excretion of hydroxyproline (15, 73, 93, 102, 116, 118, 131, 172).

Hydroxyproline is an iminoacid with a molar weight of 131.4; its rough formula is $C_5H_9O_3N$, whose structural configuration represents a 4-

hydroxy-pyrrolidine-2-carboxylic acid. Discovered by Fischer in 1902 (42, 43), in the hydrolysis products of gelatin, it was isolated in the peptidic form in urine by Westall (172).

Hydroxyproline is present in the organism to a significant extent (13%) in the collagenous stratus, which as we know, constitutes about 1/3 of the body weight (125).

The iminoacid present in the collagen is not of direct exogenous, alimentary derivation; instead it comes from the hydroxylation of the proline such as it occurs through the conversion of lysin into hydroxyline (47, 62, 90, 106, 130, 138, 158, 164).

Many interpretative hypotheses have been set up as to the time and the way in which this process of hydroxylation takes place (25, 56, 59, 61, 62, 75, 113, 115, 119, 146, 163). Regardless of the metabolic phases of this conversion, it is certain that the transition from proline into hydroxyproline is considered a direct expression of the "turnover" of the collagenous tissue (48). The urinary peptides with hydroxyprolinic content as demonstrated by recent studies with proline C₁₄, therefore assume the role

of indicators of this metabolic activity (56, 57, 134, 137, 176, 177, 186), while the rate of free iminoacid in the urine turns out to be rather small and insignificant (93, 118, 139, 86). Investigations conducted by various research groups in this connection revealed the relationships between hydroxyproline and the polypeptidic subunits which make up the collagen (89, 93, 114, 115, 118, 139, 186).

We know very well today that the first steps of collagen synthesis take place in the fibroblast (3, 44, 45, 51, 55, 58, 145): this is where the activation of the aminoacids takes place (113). The proline is hydroxylated into hydroxyproline and the lysin is converted into hydroxyline. Peptidic subunits up to 250 aminoacid, with a molar weight of about 30,000, accumulate in the ribosomes. A carbohydrate -- with an as yet unknown composition -- is added to the terminal residues of the aspartic acid, at the end of the peptide, and constitutes the link between the two peptidic chains (52). This is how the "alpha" chains are formed; each of them contains four of these peptides and has a molar weight of 120,000. Three "alpha" chains provided with weak bonds are then added to bring about the typical spiral structure of the tropocollagen with a molar weight of 360,000 (36, 37, 66, 68, 129, 153). The latter represents the fundamental complex of the collagen ("the basic building block") (147). Depending upon the progressive complexity of the intramolecular bond, three types of tropocollagen were identified: alpha, beta, and gamma (151, 154). In this phase, the tropocollagen is extracellular and it is soluble in neutral saline solution (154). By virtue of the strong bonds between contiguous chains, the tropocollagen successively loses its solubility characteristics and undergoes fibrillary metamorphosis. The massive stabilization of the fibrils and the constitution of the mature collagen would thus take place through covalent interfibrillary and intermolecular bonds between adjacent units of tropocollagen (52, 53, 54).

It has been suggested that a weak bond between the hydroxylic group of the hydroxyproline and the carboxylic group of a nearby chain could stabilize the collagenous fiber, in a manner similar to what occurs in the case of the disulfide bond of keratin (69, 70, 71). It is a fact that, as the molecular structure becomes increasingly complicated, the tropocollagen rates decrease, the solubility of the collagens is reduced, and the hydroxyprolinuria therefore shrinks (5, 14, 66, 89, 91, 154, 169, 186).

The presence of soluble collagen during the active growth phases has been traced to the more lively synthesis of the tropocollagen which would occur in such a quantity as to overcome the polymerization capacity of the

fibrils (66, 68). The metabolic inertia of the mass of insoluble collagen is well known indeed (167): the average life of the mature fiber is about 300 days (123), whereas that of "alpha" tropocollagen is about 5-24 hours (86, 89). In the normal adult, the collagen turnover would thus appear to be rather limited, which would indicate a "breakdown" under some particular conditions, that is, in the course of bone reformation [reshaping] and in the postpartum uterine involution (74, 122, 174). The catabolism of the insoluble collagen however would involve its conversion into tropocollagen and its successive hydrolysis into peptides and amino acids (84, 174).

On the basis of recent experimental investigations, it has been possible to ascertain that hydroxyprolinuria is sustained by the transition in the urine, of the imino acid into the state of metabolic precursor (active imino acid) or the state of catabolite of the collagen fiber or, likewise, of nonmetabolized alimentary hydroxyproline. The increase in hydroxyprolinuria would therefore be brought about by the increase in the anabolism of the collagen, by the stoppage of its polymerization (which occurs in osteolathyrism), or finally by increased tissue catabolism, as was also observed in extensive burns (48, 89, 98, 173). On the other hand, a reduction in hydroxyprolinuria would indicate a slowdown in the collagen metabolism due to a decrease in the processes of biosynthesis or perhaps due to an abnormal accumulation of insoluble collagens. Decreased excretion of imino acid would occur also because of a scurvy-producing diet (94, 114).

Many observations therefore seem to suggest that hydroxyprolinuria constitutes an indication proving the rate of metabolically-active collagen although in the current situation we cannot yet state specifically whether catabolic processes prevail or whether phenomena pertaining to the biosynthesis of the connective tissue predominate (48). A number of physiopathological conditions are very significant in this connection. In particular it has been observed that young animals, with a higher rate of soluble collagen and with an accelerated metabolic turnover, reveal increased hydroxyprolinuria (67, 114, 123, 124). It was also found that, in scurvy, where both the soluble collagen rate and the connective tissue synthesis decrease, a diminution of hydroxyprolinuria takes place (121). Finally it was observed that, in lathyrism, the dramatic increase in the soluble collagen is accompanied by a high urinary excretion of free and peptidic hydroxyproline. In this last-named condition, fibrillogenesis is in fact blocked in the tropocollagen stage (89, 105, 114, 160, 162).

The influence on hydroxyprolinuria, explained by hormones with osteo-metabolic action, such as the growth hormone and parathormone has been amply documented by various researchers (Bates, Dull, Keiser, Klein, Lee, Prockop, Smiley); it was thus possible to ascertain the existence of a significant relationship between urinary excretion of hydroxyproline and the turnover of tropocollagen (8, 32, 33, 92, 96, 97, 102, 144). Consequently, the study of the urinary excretion of hydroxyproline very recently turned out to be a new parameter in the evaluation of the turnover of the connective tissues during growth and in some morbid conditions (32, 33, 35, 88,

89, 96, 98, 103, 107, 116, 117, 118, 143, 149, 150, 155, 159, 186).

In our research here, we have come to feel that an examination of the behavior of hydroxyprolinuria is worth-while; we conducted this study in a series of patients whose altered endocrinometabolic situation had produced some conditions capable of affecting hydroxyprolinuria to an appreciable extent. In particular, we considered the relationships between the urinary excretion of hydroxyproline and the concentration of circulating HGH in some patients with metabolic disorders of the diabetic type.

Material and Methods

The investigations were conducted on a total of 71 subjects. In particular, we examined 40 patients with diabetes mellitus, 11 patients with acromegaly, and 20 subjects who were clinically healthy.

The age of the diabetics, who were members of both sexes, was between 9 and 78 years; in 11 cases, the metabolic syndrome developed before the age of 20; in 4 patients, it assumed the clinical and endocrinometabolic characteristics of diabetes with hypophyseary overtones.

All subjects examined were kept on a diet devoid of gelatin, fish or meat during the 24 hours prior to the time the urine samples were taken and throughout the entire time of the experiment. The urine samples were kept at a temperature of $+4^{\circ}\text{C}$ under a layer of toluene.

The blood sample for the determination of the HGH was taken at base conditions on the second day of the experiment, about 8-10 hours after the last food intake.

In all patients subjected to insulin treatment, throughout the entire period of the experiment, administration of insular hormone was suspended.

To determine the plasmatic HGH rate, we used the radioimmunity method proposed by Hunter and Greenwood (80) and modified by Berson and associates (11), with some additional features suggested by us (156).

The determination of the total hydroxyproline rate in the urine was made according to the technique proposed by Prockop and Udenfriend (140) which employs the evaluation of a chromophore with an absorption spectrum on the wavelength of 360 m μ .

Results

The results of the determinations of hydroxyprolinuria and of the circulating HGH rate are given in Tables 1, 2, and 3.

Table 1. Normal Subjects

a. N.	b. Name	c. Age	HCN mg/ml	Hydroxyproline - urinary mg/24h
1	S.N.	10	6.5	46
2	D.G.L.	9	7.5	52
3	V.R.	13	6.0	30
4	S.C.	11	5.0	42
5	C.M.	15	4.4	32
6	L.P.	12	4.3	30
7	S.A.	15	3.8	44
8	T.P.	13	5.6	38
9	M.O.	21	3.8	22
10	T.K.	20	4.1	28
11	D.M.	27	2.4	21
12	D.F.A.	20	2.0	16
13	A.C.	20	3.5	18
14	B.D.	26	2.5	15
15	C.N.	23	1.8	20
16	S.V.	19	2.9	23
17	P.F.	22	2.3	20
18	R.L.	21	4.0	14
19	G.T.	20	2.1	10
20	M.M.	20	2.0	16

Key: a. Number

b. Name

c. Age

d. Urinary hydroxyproline, mg/24 hr

The urinary excretion of hydroxyproline in normal subjects was between 10 and 52 mg/24 hr, with an average of 28.35 mg/24 hrs.

The hydroxyprolinuria registered in young subjects appeared greater than that registered in adults. Indeed, the daily elimination of imino acid varied between 30 and 52 mg/24 hours. These data agree with the observations made earlier by other researchers.

Table II. Diabetes Mellitus

a N.	b Name	c Age	d Inicio		g Intensidad del diabete	h Acidosis	i Complica- ciones	HGH mIU/ml	j Idroxi- prolina urinaria mg/24h
			Primero 22 a.	Despues 22 a.					
1	N.S.	27	+		+++	+	+	7.0	60
2	S.L.	12	+		+++	+		10.3	125
3	M.G.	9	+		+++	+	+	7.0	45
4	F.A.	13	+		+++	+		12.6	96
5	M.P.	17	+		+++	+		9.7	105
6	T.P.	13	+		+++	+		8.9	85
7	M.F.	26	+		+++	+	+	11.5	65
8	P.B.	24	+		+++			3.4	26
9	T.C.	41	+		+++		+	7.1	95
10	P.L.	28	+		++	+	+	6.8	120
11	S.I.	33	+		+			1.0	52
12	A.A.	43		+	+++	+	+	8.3	48
13	L.M.	55		+	+++	+	+	10.0	47
14	B.Z.	53		+	+++		+	4.0	52
15	S.A.	53		+	+++	+		12.0	26
16	T.D.	51		+	+++	+		8.3	28
17	C.M.	51		+	+++	+	+	5.0	80
18	E.A.	44		+	+++			1.2	33
19	M.P.	40		+	+++	+		2.0	11
20	C.A.	39		+	+++			3.3	40
21	S.A.	66		+	++		+	4.3	40
22	T.O.	53		+	++			4.0	70
23	P.P.	47		+	++		+	4.1	70
24	C.C.	50		+	++			4.3	13
25	P.A.	44		+	++			3.5	19
26	B.R.	70		+	++		+	4.8	34
27	P.E.	63		+	++		+	2.9	15
28	C.M.T.	45		+	++		+	3.4	16
29	P.A.	70		+	+			4.0	25
30	V.A.	26		+	+			5.0	20
31	D.F.	54		+	+		+	1.3	14
32	P.R.	50		+	+			3.8	18
33	S.A.	47		+	+			3.5	10
34	B.R.	35		+	+			9.6	42
35	M.O.	41		+	+			8.3	24
36	B.C.	51		+	+			1.3	50
37	C.A.	29		+	+		normog.	10.0	110
38	N.T.	57		+	+		normog.	4.8	200
39	P.E.	40		+	+		normog.	12.5	225
40	G.V.	30		+	++		normog.	12.3	225

Key: a. Number

b. Name

c. Age

d. Start

e. Before age 22

f. After age 22

g. Intensity of diabetes

h. Acidosis

i. Complications

j. Urinary hydroxy-
proline, mg/24 hrs

Table III. Acronegaly

Key: a. Number
b. Name
c. Age

d. Diabetes
e. Urinary hydroxyproline, mg/24 hrs

As regards the circulating HGH rate in the controls, this came out in the range of the norm and produced the highest values in growing youngsters (3.8-7.8 mmug/ml).

In diabetic subjects, the excretion of imino acid appeared greater than that registered in the controls, with an average value of 66.07 mg/24 hrs and a percentage increase of 133.15 percent (Figures 1 and 2). But if we consider the two groups of patients examined separately, then we can make some observations that are rather interesting. As a matter of fact, average hydroxyprolinuria turned out to be 79.45 mg/24 hrs in young diabetics and 61.0 mg in old diabetics: compared to the controls, this gave us a respective increase of 180.21 percent and 115.16 percent (Figures 2, 3, and 4). Hydroxyprolinuria thus turned out to be noticeably higher in young diabetics. On the other hand, the urinary excretion of hydroxyproline appeared particularly distinct in subjects where the diabetic syndrome was combined with clinical and humoral signs of increased somatotrophic secretion and in some acromegaly patients where metabolic disorders of the diabetic type could not be documented.

It thus appears evident that hydroxyprolinuria reveals the highest values in the hypophyseary diabetic and in the young diabetic, that is, in cases where we often have a clinical picture of most severe diabetes, a picture that is sometimes unstable and that easily involves ketoacidosis.

The fact is that the average values recorded in these two groups of patients were 227.5 mg/24 hrs in hypophyseal diabetes and 79.45 mg/24 hrs in youthful diabetes.

The concentration of HGH explored in a fasting state, under basic conditions, produced the highest values in the acromegaly patients and in some young diabetics. This last observation, reported earlier by Conti, and associates, has been confirmed once again in this series of investigations so that we have good reason to state that youthful diabetes--particularly if it is very severe--is associated quite frequently with HGH concentrations above normal.

As regards the relations between the two endocrinometabolic parameters studied in this investigation (Tables 2 and 3; Figures 2, 3, 4, and 5), we must emphasize that, in subjects with more pronounced hydroxyprolinuria, the circulating HGH concentration is often higher than the norm. This phenomenon is most evident in young diabetics (Figure 4) and in acromegaly patients with apparent diabetes: in this last group of patients, the daily excretion of hydroxyproline reaches the highest values and even the circulating HGH concentration appears exceedingly pronounced (Figure 5). However, while there is no doubt that correlations do exist between hydroxyprolinuria and the HGH rate, it must be pointed out very specifically that a strict and direct proportionality is not always recognizable between these two parameters. With equal circulating HGH concentration, we in fact observe that the hydroxyprolinuria is relatively higher in acromegaly subjects with diabetes than in young diabetics. This is obviously due to the intervention of other, not well known factors which further aggravate the protein exchange disorders in the disease of acromegaly.

Conclusions

The documented influence of hormones with osteometabolic action, such as HGH and parathormone, on hydroxyprolinuria caused us to consider bone collagen as the major source of hydroxyproline eliminated through the urine (8, 32, 33, 92, 96, 97, 102, 144). The relationships recorded between the soluble collagen rate and the urinary hydroxyproline seem to support the existence of a direct correlation between growth and hydroxyprolinuria. Gross, as a matter of fact, was able to demonstrate the existence of a precise relationship between the growth rate and the dimension of the soluble pool of collagen (66). Systematic research conducted by Jones and associates on subjects between the ages of 5 and 49 years demonstrated that hydroxyprolinuria increased from infancy to puberty; it reaches maximum values at the age of 14-15, with the excretion of 350-390 mg/24 hrs. After the age of 23, the rate of urinary excretion of hydroxyproline becomes stabilized around 30 mg/24 hrs, with daily variations of less than 10 percent. The research group of Ziff and associates was likewise able to determine an increased elimination of hydroxyproline in children in the growth phase and furthermore noted that the supply of HGH is accompanied by a definite increase in hydroxyprolinuria (72, 88, 186).

The relationships between the growth hormone and the deposit of collagen have been investigated for quite some time and the research conducted with Proline-C₁₄ documented the way in which the capture of this imino acid, by the collagen, increases as a result of the supply of somatotrophic hormone (4, 7, 26, 99, 142). As for the rest, it is well known that the important task of the growth hormone in the metabolic dynamics of the mucopolysaccharides suggested the perfection of a biological procedure for the determination of the GH (19, 20, 28, 50, 120).

In patients with acromegaly in the active phase, increased hydroxyprolinuria was found (9, 10, 31, 33, 80, 88, 168), whereas in nanosmia cases, the urinary elimination of imino acid turned out to be particularly small (33, 72, 81, 108, 132, 170). In a series of studies of various endocrine diseases, Benoit and associates, as well as Bonadonna and associates, were able to demonstrate that the excretion of hydroxyproline increases after stimulation with somatotrophic hormone, gonadotropin and thyroid hormone; the effect resulting from parathormone turned out to be inconstant (9, 10, 13). These same authors investigated the urinary excretion of hydroxyproline in 12 diabetic patients between the ages of 18 and 62. Only in one case was hydroxyprolinuria considerably increased whereas in another observation it differed little from the high limits of the norm. It must be pointed out that, in both cases, diabetes had developed at an early age and that the patients examined were subjected to insulin treatment. This last disclosure is particularly important if we consider the well-known action of insulin upon the regulation of protein exchange in general and upon the utilization of the amino acids (100, 109, 111, 112, 157, 158, 179).

On the other hand, the pathogenic importance of GH in diabetes mellitus has for a long time been considered quite seriously and has been quite validly confirmed in experimental studies and clinical observations (1, 18, 24, 23, 29, 30, 40, 41, 76, 78, 81, 82, 110, 127, 141, 171, 180, 185).

In this connection, the dosage of somatotrophic hormone in the plasma has led to interesting discoveries. Conti and associates (21, 22, 23) were able to show that the disorders in somatotrophic secretion are not foreign to endocrine disorders produced by an absolute or relative insulin deficiency. In particular, the concentration of circulating GH appeared significantly increased in youthful diabetes, especially when signs of incipient disturbance of the acid-base equilibrium are manifest. These observations, in agreement with the first discoveries of Forsman and Gerszell (46), were successively confirmed by Erlich and Randle (38, 39), Pfeiffer (123), and Greenwood (63, 65).

As regards the results obtained in this, our first preliminary research, we found that in patients with diabetes mellitus the urinary excretion of hydroxyproline appeared to be greater than the average recorded among the controls; this increase is more marked in youthful diabetes, in other words, where the metabolic disorder was more severe and unstable. On the

other hand, in patients whose acromegaly syndrome was complicated by obvious disorders in glycidic exchange, the excretion of amino acid appeared particularly pronounced: it was much greater than the--albeit very high--values that are typical of acromegaly not connected with diabetes mellitus. These discoveries become even more significant when we consider that, in the adult diabetic, the rate of hydroxyprolinuria turned out to be moderately higher than the average levels.

It seems to us therefore that we are justified in agreeing that metabolic disorders, in diabetes, also involve the collagen and that they are increasingly manifest--particularly at an early age--when the turnover of the fundamental substance is already physiologically increased with respect to the active body growth.

Our discoveries thus indicate that there is a significant relationship between hydroxyprolinuria and body growth. The fact is that in the young control subjects, in the young diabetics, and in acromegaly patients, in whom the HGH levels appeared highest, we found a noticeable increase in the urinary excretion of hydroxyproline.

It was demonstrated quite some time ago that the plasmatic levels of HGH are usually high in the youthful diabetes patient and it was assumed that hypersomatotropinism, in this morbid condition, was the expression of an early functional impairment of the hypophysis or rather the result of an incretory action of the secondary hypophysis to metabolic disorders, particularly to the limited utilization of glucose produced on the cell level by the insulin shortage. But, regardless of the underlying physiopathogenic significance of hypersomatotropinism in youthful diabetes, there is no doubt that this is one of the factors most actively involved in the cause for the increase in hydroxyprolinuria precisely because of this morbid condition.

From a general viewpoint, therefore, we can conclude that both diabetes mellitus and acromegaly represent two morbid conditions which definitely have an effect on the metabolism of collagen, to the point where they often trigger a distinct increase in hydroxyprolinuria. The metabolic disorder furthermore seems to become more severe when the two sickness pictures are combined so as to give rise to acromegalic diabetes.

This conclusion, in our opinion, cannot be challenged by the observation that, in some cases of diabetes and acromegaly, there is normal hydroxyprolinuria; there are many factors which, though they are not a part of the disease of diabetes and acromegaly, do become involved in the intimate economy of collagen.

Some authors, particularly Benoit and associates, as well as Bonadonna and associates, deny that there are any noticeable variations in hydroxyprolinuria in diabetes mellitus: but in this connection we want to point out that both groups of researchers conducted their investigations

primarily on adult diabetics who were kept under control with insulin, in other words, in subjects who were in a situation definitely different from that of the patients whom we examined.

The relationships between circulating HGH and hydroxyprolinuria appear to be somewhat more complex when we compare the results obtained in some young diabetics and in acromegaly patients with diabetes. In these two groups of patients, as a matter of fact, we cannot always recognize a strict correlation between the level of somatotrope and the volume of hydroxyprolinuria because, with an equal concentration of circulating GH, the maximum excretion of hydroxyproline is characteristic of acromegaly with diabetes. This obviously documents the fact that insulin shortage aggravates the metabolic disorder of collagen to the point where it promotes a subsequent noticeable increase in the urinary excretion of imino acid. We must not be astonished that, in acromegaly without any connected apparent disorders in glycidic exchange, the behavior of hydroxyprolinuria is often changeable and not strictly correlated with the level of circulating HGH: this fact likewise figures in the easy variation of the processes of bone reshaping, through the intervention of those rather disparate factors which are present during the deposit and reabsorption of collagen.

Summary

The relationships between the concentration of circulating HGH and the urinary excretion of hydroxyproline were studied in 71 subjects, including 40 with diabetes mellitus, 11 with acromegaly, and 20 who were clinically healthy.

The plasmatic HGH was dosed with the radioimmuno-electrophoretic method of Hunger and Greenwood and the determination of total urinary hydroxyproline was made according to the technique of Prockop and Udenfriend.

It was demonstrated that in patients with diabetes mellitus the urinary excretion of hydroxyproline is often greater than the average figures recorded in the controls. This increase is more marked in youthful diabetes, in other words, where the metabolic disorder is more severe and unstable. On the other hand, in patients in whom the acromegaly syndrome was complicated by obvious disorders in glycidic exchange, the excretion of the imino acid appears particularly pronounced: it is very much greater than the--by the way in themselves rather high--values that are typical of acromegaly not connected with diabetes mellitus.

It was found that there is a significant relationship between hydroxyprolinuria and body growth. Indeed, in the controls, in young diabetics, and in acromegaly patients, in whom the HGH levels are highest, we regularly find a noticeable increase in the urinary excretion of hydroxyproline.

However we did not always find a strict correlation between the plasmatic level of the somatotrophic hormone and the urinary excretion of imino

acid, particularly in young diabetics in acromegaly patients with diabetes because--with an equal concentration of somatotrophic hormone--the higher values of hydroxyprolinuria appeared characteristic of acromegaly with diabetes.

This discovery--according to the authors--documents the fact that, in diabetes mellitus, insulin shortage helps aggravate the metabolic disorder of the collagen.

It is concluded that diabetes mellitus and acromegaly are two morbid conditions which have a profound effect on collagen metabolism and that the metabolic disorder is aggravated when the two disease pictures are combined into acromegalic diabetes.

Bibliography

1. Atkinson, R.R.B., "Acromegaly," J Boll, Son and Danielson, Ltd, London, 1932.
2. Almquist, S., "Studies on Sulfation Factor Activity of Human Serum Stockholm," Zetterlund and thelanders Boktryckeri AB, 1961.
3. Amprino, R., "Autoradiographic Research on the S 35 Sulphate Metabolism in Cartilage and Bone Differentiation and Growth, Acta Anat, 24, 121, 1955.
4. Asling, C.W., Simpson, M.G., Mook, H.D., Li C. H., Evans, H.M., "Hypophyseal Growth Hormone Nature and Actions" R.W. Smith, O.H. Graebler, C.N.H. Long, Ed, N.Y., McGraw Hill, p 576, 1955.
5. Bakerman, S., "Quantitative Extraction of Acid Soluble Human Skin Collagen with Age" Nature, 196, 375, 1962.
6. Bal'kov, B., Castello, R.A., Lozano, E.R. "Urinary Peptide Excretion by the Burned Patient," Clin Chem, 3, 217, 1957.
7. Banfield, W.G., "Effect of GH on Acetic Acid Extractable Collagen of Hamster Skin," Proc Soc Exp Biol Med, 97, 309, 1958.
8. Bates, W.K., McGowan J., Talmager, V., "Influence of the Parathyroids on Plasma Hydroxyproline Levels," Endocrinology, 71, 189, 1962.
9. Benoit, F.C., Theil, G.B.- Watten, R.H., "Hydroxyproline Excretion in Endocrine Disease," Metabolism, 12, 1072, 1963.
10. Benoit, F.L., Theil, G.B., Watten, R.H., "Urinary Hydroxyproline Excretion in Endocrine Disease," Clin Res, 11, 89, 1963.

11. Berson, S.A., Yalow, R.S., "Immunoassay of GH in Serum," 45th Meet Endocrine Soc, Atlantic City, June 1963.
12. Biserte, G., Breton, A., Fontaine, G., "Analytical Study of Urinary Amino Acids in Common Childhood Rachitism," Arch Franc Pediat, [French Pediatrics Archives], 12, 988, 1955.
13. Bonadonna, G., Sonnenberg, M., Merlino, M.I., "Total Urinary Hydroxyproline Excretion in Diabetics Before and After Hypophysectomy and After GH in Adult Hypopituitarism," Metabolism, 14, 832, 1965.
14. Boucek, R.I., Noble, N.L. Woessner, J.F. "The Effects of Tissue Age and Sex Upon Connective Tissue Metabolism," Ann N.Y. Acad Sci, 72, 1016, 1959.
15. Boulanger, P., Biserte, G., "Free and Combined Amino acids in Human Urine," Exposés Ann Biochim Med [Medical Biochemistry Yearbook Report], 18, 123, 1956.
16. Boulanger, P., Biserte, G., Courtot, F., "Paper Chromatography of Amino Acids and Polypeptides in Biological Liquids; 3 Aminoacids and urinary peptides," Bull Soc Chim Biol [Bulletin of the Chemical and Biological Society], 34, 366, 1952.
17. Carsen, M.E.A., "A Procedure for the Ion Exchange Chromatography of the Amino Acids of Urines," J. Am. Chem Soc, 74, 5954, 1952.
18. Coggeshall, C., Root, H.F., "Acromegaly and Diabetes Mellitus," Endocrinology, 26, 1, 1940.
19. Collins, E.J., Baker, V.F., GH and Radiosulfate Incorporation; a New Assay Method for GH," Metabolism, 9, 556, 1960.
20. Collins, E.J., Lysters, C., Carpenter, O.S., "Growth Hormone and Radiosulfate Incorporation in Cartilage, Comparison with the Tibia Test," Acta Endocrinol, 36, 51, 1961.
21. Conti, C., Sereno, L., "Modern Discoveries in the Field of Somatotrophic physiology and Physiopathology--Experimental and Clinical Contributions," Relaz Aiunioni Mediche-Chirurgiche Internazionali di Torino [Report of the Turin International Medical-Surgical Meetings], 1961; Folia Endocrinol, 14, 300, 1961.
22. Conti, C., Sereno, L., "The Human Somatotrophic Hormone--Experimental and Clinical Findings," VII Congr Naz Soc It Endocrinol [7th Congress of the Italian National Endocrinological Society], Pavia, 1963.
23. Conti, C., Sereno, L., "Some Contributions to the Study of the Somatotrophic Activity in Diabetes Mellitus," Intern Diabetes Fed, 5th Congress, Toronto, 1964.

24. Cushing, H., Davidoff, L., Acromegaly, Basal Metabolism, Arch Int Med, 39, 673, 1927.
25. Cushing, H., Davidoff, L., Human Acromegaly, Monograph, Rockefeller Inst Med Res, 22, 1, 1927.
26. Daughaday, W.H., Mariz, I.K., Conversion of Proline C₁₄, Labelled to Hydroxyproline by the Rat Cartilage in Vitro; Effects of Hypophysectomy, GH, and Cortisol, J. Lab Clin Med, 59, 741, 1962.
27. Daughaday, W.H., Mariz, I.K., The Formation of Free Hydroxyproline by Rat Cartilage in Vitro, J. Biol. Chem., 237, 2831, 1962.
28. Denko, C.W., Bergenstal, D.M., The Effect of Hypophysectomy and GH on Sulfur-35 in Fixation in Cartilage, Endocrinology, 57, 76, 1955.
29. Di Raimondo, C.V., Biglieri, E.C., Mitchell, D.B., Watlington, C.O., Forsham, P.H., Metabolic Analysis of HGH Activity in Man, I Congr Int Endocrinol, Copenhagen, 1960.
30. Dominguez, J.M., Greenberg, A.G., Pazianos, B.S., Pearson, O.H., Diabetogenic and Hypoglycemic Effects of HGH, 1st Congr Int Endocrinol, Copenhagen, 1960.
31. Dominguez, J.M., Pearson, O.H., Immunologic Measurement of GH in Human Sera, J. Clin Endocrinol Metab, 22, 865, 1962.
32. Dull, T.A., Causang, L., Henneman, P.H., Urinary Total Hydroxyproline as an Index of Connective Tissue Turnover in Bone, J. Clin Invest, 41, 1355, 1962.
33. Dull, T.A., Henneman, P.H., Urinary Hydroxyproline as an Index of Collagen Turnover in Bone, New England J Med, 268, 132, 1963.
34. Ebert, F.S., Prokop, D.J., The Urinary Hydroxylation of Proline to Hydroxyproline During the Synthesis of Collagen in Chick Embryos, Biochem and Biophysics Research Commun, 8, 305, 1962.
35. Efron, M.L., Bixby, E.M., Palattao, L.G., Pryles, C.V., Hydroxyprolinemia Associated with Mental Deficiency, New Engl J. Med, 267, 1193, 1962.
36. Engel, J., "On the Formation of Fibrils and Long-Spacing Segments from Denatured Tropocollagen Solutions," Z Physiol Chem [Jr of Physiological Chemistry], 329, 69, 1962.
37. Engel, J., Investigation of the Denaturation and Renaturation of Soluble Collagen by Light Scattering, Arch Biochem Biophys, 97, 150, 1962.

38. Erlich, R.M., Randle, P.J., Serum GH Concentrations in Diabetes Mellitus, *Lancet*, 2, 233, 1961.
39. Erlich, R.M., Randle, P.J., Immunoassay of Serum GH in Diabetes Mellitus, *Ciba Found Coll Endocrinol*, 14, 117, 1962.
40. Escamilla, R.F., Hutchins, J.J., Deanner, W.C., Li, C.H., Clinical Experiences with GH in Pituitary Infantilism and Gonadal Dyagenesis, 1st Congr Int Endocrinol, Copenhagen, 1960.
41. Evans, H.M., Simpson, M.E., Reichert, F.W., Disturbance of Carbohydrate Metabolism in Normal Dogs Injected with Hypophyseal GH, *Proc Soc Exp Biol Med*, 29, 857, 1932.
42. Fisher, E., quoted in *Traite de Chimique organique [Organic Chemistry Treatise]*, XIX, p 147, Masson and C., Ed, Paris, 1962.
43. Fischer, E., Bergell, P., *Ber*, 35, 3785, 1902.
44. Fitton-Jackson, S., Cytoplasmic Granules in Fibrogenic Cells, *Nature*, 175, 39, 1955.
45. Fitton-Jackson, S., Smith, R.H., Studies on the Biosynthesis of Collagen. The Growth of Fowl Osteoblasts and the Formation of Collagen in Tissue Culture, *J. Biophys Biochem Cytol*, 3, 897, 1957.
46. Forsmann, O., Gemzell, C.A., Plasma Level of GH in Patients with Diabetes Mellitus, Hypercholesteremia and Liver Diseases, *Acta Endocrinol*, 32, 430, 1959.
47. Frankland, D.M., Wynn, C.H., Degradation of Acid Soluble Collagen by Rat Liver Preparations, *Biochem J*, 85, 276, 1962.
48. Frey, M.Y., "Value of the Exploration of Hydroxyprolinemia and Hydroxyprolinuria in the Study of Collagen Metabolism Disorders," *J Med Lyon*, [*Lyon J. of Medicine*], 45, 1547, 1964.
49. Fujimoto, D., Tamiya, H., Incorporation of O-18 from Air into Hydroxyproline by Chick Embryo, *Biochem J*, 84, 333, 1962.
50. Gabrilow, J.L., Schwartz, A., Churg, J., Effect of Hormones on the Skin in Endocrinologic Disorders, 1st Clin Endocrinol Meet, 22, 688, 1962.
51. Gaines, L.M., Synthesis of Acid Mucopolysaccharides and Collagen in Tissue Cultures of Fibroblasts, *Bull Johns Hopkins Hosp*, 106, 195, 1960.
52. Gallop, P.M., Concerning Some Special Structural Features of the Collagen Molecule, *Biophys J*, 4 Suppl 79, 1964.

53. Gallop, P.M., Blumenfeld, O.O., Franzblau, G., Seifert, G., Intramolecular Ester Linkages and Other Unusual Bands in Collagen, Abstr. 141 Meet Am Chem Soc, p 9 C, 1962.
54. Gallop, P.M., Seifter, S., Heilman, E., Occurrence of "Ester Like" Linkages in Collagen, Nature, 183, 1659, 1959.
55. Gerber, A.B., Altmann, K.J., Mechanism of Collagen Synthesis, Nature, 183, 165, 1961.
56. Gerber, G.B., Gerber, G., Altman, J.K., Some Interrelated Aspects of Proline and Hydroxyproline Metabolism, Nature, 185, 767, 1960.
57. Gerber, G.B., Traelvels, K.R., Wood, D., Altman, K.I., The Quantitative Determination of Pyrrole-2-carboxylic Acid in Urine, Clin Chem, 9, 185, 1964.
58. Gerish, J., Catchpole, H.R., The Organization of Ground Substance and Basement Membrane and Its Significance in Tissue Injury, Disease and Growth, Am J Anat, 85, 457, 1949.
59. Gould, B.S., Biosynthesis of Collagen. The Direct Action of Ascorbic Acid on Hydroxyproline and Collagen Formation in Subcutaneous Polyvinyl Sponge Implants in Guinea Pigs, J. Biol. Chem., 232, 637, 1958.
60. Gould, B.S., Manner, G., Goldman, H.M., Stolman, J.N., Some Aspects of Collagen Formation, Ann N.Y. Acad Sci, 85, 385, 1960.
61. Gould, B.S., Woessner, J.F., Biosynthesis of Collagen, J. Biol Chem, 226, 289, 1961.
62. Green, N.M., Lowther, D.A., Formation of Collagen Hydroxyproline in Vitro, Biochem J, 71, 55, 1959.
63. Greenwood, F.C., A Radioimmunoassay for BCG and Its Applications, Meet Immunol. Prot. Horm., Rome, 1964.
64. Greenwood, F.C., Hunter W.M., Glover, J.S., The Preparation of ^{131}I Labelled BCG of High Specific Radioactivity, Biochem J., 89, 114, 1963.
65. Greenwood, F.C., Hunter, W.M., Marrian, V.J., GH Levels of Children and Adolescents, Brit Med J, 1, 25, 1964.
66. Gross, J., Studies on the Formation of Collagen. The Influence of Growth Rate on Neutral Salt Extracts of Guinea Pig Dermis, J. Exp Med, 107, 265, 1958.

67. Gross, J., Studies on the Formation of Collagen. Effect of Vitamin C Deficiency on the Neutral Salt Extractable Collagen of Skin, *J. Exp Med*, 109, 557, 1959.
68. Gross, J., Collagen, *Sci Am*, 204, 120, 1961.
69. Gustavson, K.H., The Chemistry and Reactivity of Collagen, Acad Press, N.Y., 1956.
70. Gustavson, K.H., Acta Chem Scand. [Scandinavian Chemical Transactions], 8, 1299, 1954.
71. Gustavson, K.H., Svensk Kem Tidskr [Swedish Chemical Journal], 54, 74, 1942.
72. Haak, A., Kassensar, A.A.H., Querido, A., in Protein, Metabolism. Influence of GH, Anabolic Steroids and Nutrition in Health and Disease, ed by Gross, F., Berlin, Springer Publishers, p 150, 1962.
73. Hansen, H., Fittkav, S., "On Peptides in the Human Urine," J Physiol Chem, 313, 152, 1958.
74. Harkness, R.D., Morales, B.E., The Time, Course, and Route of Loss of Collagen from the Rat's Uterus During Post-Partum Involution, *J. Physiol*, 132, 502, 1956.
75. Hausmann, E., Neuman, W.F., Conversion of Proline to Hydroxyproline and Its Incorporation into Collagen, *J. Biol Chem*, 149, 236, 1961.
76. Henneman, D.H. Henneman, P.H., Effects of HGH on Levels of Blood and Urinary Carbohydrate and Fat Metabolites in Man, *J. Clin Invest*, 39, 1239, 1960.
77. Henriques, V., Sorensen, S.P.L., "On the Quantitative Determination of Amino Acids, Polypeptides, and Hippuric Acid in the Urine by Means of Formol Titration," Z. Physiol Chem, 64, 120, 1909.
78. Hernberg, C.A., The Effect of HGH on Severe Juvenile Diabetes After Hypophysectomy, *Acta Endocrinol*, 33, 559, 1960.
79. Housay, B.A., Anderson, E., Diabetogenic Action of Purified Anterior Pituitary, *Endocrinology*, 45, 627, 1949.
80. Hunter, W.M., Greenwood, F.C., A Radioimmuno-electrophoretic Assay for HGH, *Biochem J*, 85, p 39, 1962.
81. Ikkos, D., Luft, R., Aspects of the Metabolic Action of HGH, *Ciba Found Coll Endocrinol*, 13, 107, 1960.

82. Ikkos, D., Luft, R., Studies on the Metabolic Action of HGH
1st Congr Inter Endocrinol, Copenhagen, 1960.
83. Ikkos, D., Luft, R., Gemzell, C.A., Almquist, S., Effect of
HGH on Glucose Tolerance and Some Intermediary Metabolites in Man,
Acta Endocrinol, 39, 547, 1962.
84. Jackson, D.S., The Formation and Breakdown of Connective Tissue.
Connective Tissue., Springfield, Illinois, Ch Thomas, 62, 1957..
85. Jackson, D.S., Metabolic Studies: Turnover of Alpha and Beta
Chains. Symposium on Molecular Biology of Collagen, Div Biol Chem, Am
Chem Soc, 21, 1962.
86. Jackson, D.S., Bentley, J.P., On the Significance of the Extra-
ctable Collagens, J. Biophys Biochem, Cytol, 7, 37, 1960.
87. Jasin, H.E., Fink, C.W., Smiley, W.S., Ziff, M., Effect of GH
on Urinary Hydroxyproline, Clin Res, 10, 34, 1962.
88. Jasin, H.E., Fink, C.W., Wise, W., Ziff, M., Relationship Be-
tween Urinary Hydroxyproline and Growth, J. Clin Invest, 41, 1923, 1962.
89. Jasin, H.E., Ziff, M., Relationship Between Soluble Collagen
and Urinary Hydroxyproline in the Lathyrotic Rat, Proc Soc Exp Biol Med,
110, 837, 1962.
90. Jones, C.R., Bergman, M.W., Kittner, R.J., Pigan, W.W., Urinary
Hydroxyproline Excretion in Normal Children and Adolescents, Proc Soc
Exp Biol Med, 115, 85, 1964.
91. Kao, K.Y.T., McGavack, T.H., Connective Tissue. Age and Sex
Influence on Protein Composition of Rat Tissues, Proc Soc Exp Biol Med,
101, 153, 1959.
92. Keiser, H.R., Gill, J.R., Sjoerdsma, A., Bartter, F.C., The
Effect of Para thyroid Extract on Hydroxyproline Metabolism, Clin Res,
11, 41, 1963.
93. Kibrick, A.C., Hashiro, C.Q., Safier, L.B., Hydroxyproline
Peptides of Urine in Arthritic Patients and Controls on a Collagen-Free
Diet, Proc Soc Exp Biol Med, 109, 473, 1962.
94. Kibrick, A.C., Hashiro, C.Q., Walters, M; Milhorat, A.T.,
Hydroxyproline Excretion in Urine of Patients with Muscular Dystrophy and
Others Muscle Diseases, Proc Soc Exp Biol Med, 115, 662, 1964.
95. Kivirikko, K.I., Liesmaa, M., Effect of Cortisone and Age on
the Hypro and Prolina Concentrations of Blood and Urine in Rat, Acta
Endocrinol, 27, 441, 1958.

96. Klein, L., Albertson, K., Curtiss, P.H., Urinary Hypro in Hyperparathyroidism: a Study of 3 Cases with and Without Bone Lesions, *Metabolism*, 11, 1023, 1962.

97. Klein, L., Curtiss, P.H., Jr., Effect of Parathyroid Extract Upon the Urinary Excretion of Peptide Hydroxyproline, *Feder Proc*, 21, 206, 1962.

98. Klein, L., Curtiss, P.H., Davis, J.H., Collagen Breakdown in Thermal Burns, *Surg Forum*, 13, 459, 1962.

99. Knobil, E., in *Growth in Living Systems*, M.X. Zarrow, N.Y., Basic Book, Inc, p 353, 1961.

100. Krah1, M.E., Function of Insulin and Other Regulatory Factors in Peptide Formation by Animal Cells, *Rec Progr in Hormone Research*, 12, 199, 1956.

101. Lampert, D.T., Hydroxyproline Biosynthesis: Loss of Hydrogen During the Hydroxylation of Proline, *Nature*, 202, 293, 1964.

102. Lee, C.A., Urinary Hypro in Diseases Involving Bone and Calcium Metabolism, *Med J. Aust*, 1, 892, 1964.

103. Lenzi, F., Ravenni, G., Rubegni, M., Del Giovane, L., "Significance of Urinary Hydroxyprolinuria in Osteoporosis," *Panminerva Med*, 5, 155, 1962.

104. Lenzi, L., Silva, E., "Current Concepts of Physiology and New Orientation in the Pathology of the Connective Tissue: Collagen," *Chir Organi Mov [Mobile Organ Surgery]*, 53, 245, 1964.

105. Levene, C.J., Gross, J., Alterations in State of Molecular Aggregation of Collagen in Chick Embryos by β -aminopropionitrile, *J. Exp Med*, 110, 771, 1959.

106. Levine, M., Conversion of Proline to Hydroxyproline in Rat Skin in Vitro, *Fed Proc*, 21, 169, 1962.

107. Lindstedt, S., Prockop, D.J., Isotopic Studies on Urinary Hypro as Evidence for Rapidly Catabolized Forms of Collagen in the Young Rat, *J. Biol Chem*, 236, 1397, 1961.

108. Lipssett, M.B., Bergenstal, D.M., Dhyse, F.B., Metabolic Studies with BCH in Dwarfism and Acromegaly, *J. Clin Endocrinol Metab*, 21, 119, 1961.

109. Lukens, F.D.W., The Influence of Insulin on Protein Metabolism, *Diabetes*, 2, 491, 1953.

110. McKerns, K.W., Discussion in Raben, M.S., Human GH, Rec Progr Horm Res, 15, 106, 1959.

111. Manchester, K.L., Young, F.G., The Effect of Insulin on Incorporation of Amino Acids into Protein of Normal Rat Diaphragm in Vitro, Biochem J., 70, 353, 1958.

112. Manchester, K.L., Young, F.G., Location of C₁₄ in Protein from Isolated Rat Diaphragm Incubated in Vitro with C₁₄ Amino Acids and with C₁₄ O₂, End Proc 18, 280, 1959.

113. Manner, G., Could, B.S., Collagen Biosynthesis: the Formation of Hypro and Soluble Ribonucleic Acid Complexes of Proline and Hypro by Chick Embryo in Vivo and by a Subcellular Chick Embryo System, Biochim Biophys Acta, 72, 243, 1963.

114. Martin, G.R., Mergenhagen, S.E., Prockop, D.J., Influence of Scurvy and Lathyrism on Hypro Excretion, Nature, 191, 1008, 1961.

115. Martin, G. R., Piaz, K.A., Lewis, M.S., The Incorporation of C₁₄ Glycine into the Subunits of Collagens from Normal and Lathyrotic Animals, Biochim Biophys Acta, 69, 472, 1963.

116. Mechanic, G., Skupp, S.J., Safier, L.B., Kibrick, A.C., Isolation of 2 Peptides Containing Hypro from Urine of a Patient with Rheumatoid Arthritis, Arch Biochem, Biophys, 86, 71, 1960.

117. Meilman, E., Urivetsky, M.M., Studies on Urinary Hypro, Arth and Rheum, 4, 119, 1961.

118. Meilman, E., Urivetsky, M.M., Rapoport, C.M., Urinary Hypro Peptides, J Clin Invest, 42, 40, 1963.

119. Maister, A., Stone, E., Manning, J.M., Conversion of Proline to Hypro in Collagen Synthesis, Abs of 142d Meet Am Chem Soc, p 19, 1962.

120. Meyer, K., in Struktur und Stoffwechsel des Bindegewebes [Structure and Metabolism of Connective Tissue], Hauss and Loose, Stuttgart, p 12, 1960.

121. Mitoma, C., Smith, I.E., Studies on the Role of Ascorbic Acid in Collagen Synthesis, J. Biol Chem, 235, 426, 1960.

122. Morrione, T.C., Seifter, S., Alteration in the Collagen Content of the Human Uterus During Pregnancy and Postpartum Involution, J. Exp Med, 115, 357, 1962.

123. Benbarber, A., Perrone, J.C., Slack, H.C.B., The Relative Metabolic Inertia of Tendon Collagen in the Rat, Biochem J, 49, 199, 1951.

124. Neuberger, A., Slack, H.G.B., The Metabolism of Collagen from Liver, Bone, Skin and Tendon in the Normal Rat, *Biochem J*, 53, 47, 1953.
125. Neuman, R.E., Hydroxyproline Content of the Developing Chick Embryo, *Proc Soc Exp Biol Med*, 76, 289, 1951.
126. Orekhovitch, V.N., Shpikiter, V.O., Mazurov, V.I., Kunina O.V., Procollagens. Classification, Metabolism, Action of Proteinases, *Bull Soc Chim Biol*, 42, 505, 1960.
127. Pearson, C.H., Dominguez, J.M., Greenberg, E., Pazianos, H., Ray, B.S., Diabetogenic and Hypoglycemic Effects of HGH, *Transact Ass Am Physicians*, 73, 213, 1960.
128. Pfeiffer, E.F., "Growth Hormone and Insulin Secretion," 2d Symp Deutsch Ges fuer Endokrinologie [2nd Symposium of the German Endocrinology Society], Duesseldorf, p 41, 1964.
129. Piez, K.A., Eigner, E.A., Lewis, M.S., The Chromatographic Separation of Amino Acids and Compositions of the Subunits of Several Collagens, *Biochemistry*, 2, 58, 1963.
130. Piez, K.A., Likins, R.C., The Conversion of Lysine to Hydroxylysine and Its Relations to the Biosynthesis of Collagen, *J. Biol Chem*, 229, 101, 1957.
131. Flaguët, P., Biserte, G., Boulanger, P., "Amino Acids and Polypeptides of Biological Environments: Combined Forms of Proline, Hypro and Glycocol of Human Urine, *Bull Soc Chim Biol*, 42, 393, 1960.
132. Prader, A., Illig, R., Influence of Growth Hormone, Anabolic Steroids and Nutrition in Health and Disease, in: *Protein, Metabolism*, Gross, p 383, Berlin 1962.
133. Prockop, D.J., Collagen Degradation and Urinary Hypro, *Fed Proc* 21, 169, 1962.
134. Prockop, D.J., Isotopic Studies on Collagen Degradation and the Urine Excretion of Hypro, *J. Clin Invest*, 43, 453, 1964.
135. Prockop, D.J., Studies with Proline- H_3 on the Hydroxylation of Proline During Collagen Synthesis in Chick Embryos, *Arch Biochem*, 106, 112, 1964.
136. Prockop, D.J., Kaplan, A., Udenfriend, S., Oxygen 18: Studies on the Conversion of Proline to Hypro, *Biochem, Biophys. Research Commun*, 9, 162, 1962.
137. Prockop, D.J., Kaiser, H.R., Sjoerdsma, A., Gastrointestinal Absorption and Renal Excretion of Hypro Peptides, *Lancet*, 2, 527, 1962.

- 5
138. Prockop, D.J., Mitoma, C., Sjoerdama, A., Observations on Hypro Metabolism in Man, *Fed Proc*, 18, 434, 1959.
 139. Prockop, D.J., Sjoerdama, A., Significance of Urinary Hypro in Man, *J. Clin Invest*, 40, 843, 1961.
 140. Prockop, D.J., Udenfriend, S., A Specific Method for the Analysis of Hypro in the Tissues and Urine, *Analyt Biochem*, 1, 228, 1960.
 141. Raben, M.S., Growth Hormone, *New Engl J Med*, 266, 31, 1962.
 142. Raben, M.S., Minton, P.R., Mitchell, M.L., in *Protein Metabolism*, Cross-Berlin, Springer Publishers, p 121, 1962.
 143. Ravenni, G., Rubegni, M., Del Giovane, L., Hydroxyprolinuria in Normal Persons, *Boll Soc Ital Biol Sper*, 38, 263, 1962.
 144. Rasmussen, H., Parathyroid Hormone: Nature and Mechanism of Action, *Am J Med*, 39, 112, 1961.
 145. Robbins, W.C., Warson, R.F., Peppas, C.D., Porter, K.R., Some Effects of Anti-Collagen Serum on Collagen Formation in Tissue Culture, *J Biophys Biochem Cytol*, 1, 381, 1955.
 146. Robertson, W., The Biochemical Role of Ascorbic Acid in Connective Tissue, *Ann N.Y. Acad Sci*, 92, 159, 1961.
 147. Robertson, W., Metabolism of Collagen in Mammalian Tissues, *Biophys J*, 4, Suppl 93, 1964.
 148. Robertson, W., Hewitt, J., Augmentation of Collagen Synthesis by Ascorbic Acid in Vitro, *Biochim Biophys Acta*, 49, 404, 1961.
 149. Rodnan, G.P., Cammarata, R.J., Urinary Excretion of Hypro in Progressive Systemic Sclerosis, *Clin Res*, 11, 179, 1963.
 150. Rubegni, M., Ravenni, G., Del Giovane, L., Increased Excretion of Hypro in Rheumatoid Arthritis, *Boll Soc Ital Biol Sper*, 38, 879, 1962.
 151. Rubin, A.L., Pfahl, D., Speakman, P.T., Davison, P.F., Schmitt, F.O., Tropocollagen: Significance of Protease Induced Alterations, *Science*, 139, 37, 1963.
 152. Sarnaacka-Keller, M., The Peptide Structure of the Prothelic Acids, *Acta Biochim Polon*, 8, 383, 1961.
 153. Schleyer, M., "Separation of Alpha and Beta Components from Tropocollagen: Amino Acid Sequences of Collagens," *V.Z. Physiol Chem* 329, 97, 1962.

154. Schmitt, F.O., "Biochemical Processes in Tropocollagen," Das Leder [Leather], 14, 1, 1963.
155. Seakins, J.W.T., Peptiduria in an Unusual Bone Disorder, Biochem J, 84, 115, 1962.
156. Sereno, L., Isidori, A., Recchia, O., "First Observations on the Dosage of HGH with the Radioimmunity Method," Folia Endocrinol, 17, 598, 1964.
157. Sinex, F.M., McMullen, J., Hastings, A.B., "The Effect of Insulin on the Incorporation of C_{14} into the Protein of Rat Diaphragm," J. Biol Chem, 198, 615, 1952.
158. Sinex, F.M., Van Slyke, D.D., Christman, D.R., "The Source and State of Hydroxylisine of Collagen," J. Biol Chem, 234, 918, 1959.
159. Sjoerdsma, A., Davidson, J.D., Udenfriend, S., Mitoma, C., "Increased Excretion of Hypro in Marfan's Syndrome," Lancet, 2, 994, 1958.
160. Smiley, J.D., Yeager, H., Ziff, M., "Collagen Metabolism in Osteolathyrism: Site of Action of B-Amino Propionitrile," J. Exp Med, 116, 45, 1962.
161. Smiley, J.D., Ziff, M., "Urinary Hypro Excretion and Growth," Physiological Review, 44, 30, 1964.
162. Smith, D.J., Shuster, R.C., "Biochemistry of Lathyrism: Collagen Biosynthesis in Normal and Lathyrotic Chick Embryos," Arch Biochem Biophys, 98, 498, 1962.
163. Smith, R.H., "The Biosynthesis of Some Connective Tissue Components," Progr Biophysics and Biophys Chem, 8, 217, 1957.
164. Stetten, M.R., "Some Aspects of the Metabolism of Hypro Studied with Isotopic Nitrogen," J. Biol Chem, 181, 31, 1949.
165. Stetten, M.R., "Metabolic Relationship Between Glutamic Acid Proline, Hypro Ornithine, Amino Acids Metabolism," Baltimore, J. Hopkins Press, p 277, 1955.
166. Stone, N., Meister, A., "Function of Ascorbic Acid in the Conversion of Proline to Collagen Hypro," Nature, 194, 555, 1962.
167. Thompson, R.C., Ballov, J.E., "Studies of Metabolic Turnover with Tritium as a Tracer," J. Biol Chem, 223, 795, 1956.

163. Utiger, R.D., Parker, M.L., Daughaday, W.H., Studies on HGH: a Radioimmunoassay for HGH, J. Clin Invest, 41, 254, 1962.
169. Versar, F., Willenagter, H., "The Aging of Collagen in the Skin and in Scars," Schweiz Med Wochschr, 91, 1234, 1961.
170. Vest, M., Girard, J., "Metabolism Effects of Purified Human Growth Hormone in Dwarf Growth and in Anorexia Nervosa in Childhood," Deutsch Med Wschr, 87, 1705, 1962.
171. Vest, M., Widner, R., Kurtsteiner, W., Schmid, R., The Influence of HGH on the Metabolism and Growth in Hypopituitary Dwarfism, Schweiz Med Wschr, 90, 1468, 1960.
172. Westall, R.G., The Amino Acids Other Ampholytes of Urine, Biochem J, 60, 247, 1955.
173. Wiss, O., On Microdetermination of Oxyproline in Blood and Urine, Helv Chim Acta, 32, 149, 1949.
174. Woessner, J.F., Catabolism of Collagen and Noncollagen Protein in the Rat Uterus During Postpartum Involution, Biochem J, 83, 304, 1962.
175. Woessner, J.F., Collagen Degradation in Involuting Uterus, Fed Proc, 18, 461, 1959.
176. Woessner, J.F., Noble, N.L., Boucek, R.J., Synthesis and Degradation of Collagen in the Rat, Ex Ann Bioch Med, 24, 203, 1963.
177. Wolf, G., Berger, C.R., The Metabolism of Hypro in the Intact Rat, Incorporation of Hypro into Protein and Urinary Metabolites, J. Biol Chem, 230, 231, 1958.
178. Woodson, H.W., Hier, S.W., Salomon, J.D. Urinary Excretion of Amino Acids, by Human Subjects on Normal Diets, J. Biol Chem, 172, 613, 1948.
179. Wool, J.C., Effect of Insulin on Accumulation of Radioactivity from Amino Acids by Isolated Intact Rat Diaphragm, Nature, 202, 1967, 1964.
180. Young F.G., The Relation of the Anterior Pituitary to Carbohydrate Metabolism, Brit Med J, 11, 393, 1939.
181. Young F.G., Growth and Diabetes in Normal Animals Treated with Pituitary Diabetogenic Extracts, Biochem J, 39, 515, 1945.
182. Young F.G., Experimental Diabetes Mellitus, Schweiz Med Wochr, 76, 894, 1946.

183. Young F.G., The Experimental Approach to the Problem of Diabetes Mellitus, Brit Med J, 2, 1167, 1951.

184. Young F.G., GH and Diabetes, Rac Progr Horm Res, 8, 471, 1953.

185. Young F.G., Korner, A., Growth Hormone, in: Diabetes, Williams Ed, N.Y. 1960.

186. Ziff, M., Kibrick, A., Dresner, E., Gribetz, H.J., Excretion of Hydroxyproline in Patients with Rheumatic and Nonrheumatic Diseases, J Clin Invest, 35, 579, 1956.

FIGURE APPENDIX



Figure 1.

Key: a. Normal subjects

b. Urinary hydroxyproline mg/24 hrs




Figure 2. 1




Figure 3
Key: a. Adult diabetics
b. Urinary hydroxyproline, mg/24 hrs




Figure 4. Key: a. Youthful diabetes
b. Urinary hydroxyproline, mg/24 hrs




Figure 3.

- Key: a. Acromegaly patients
 b. Acromegaly patients with diabetes
 c. Urinary hydroxyproline, mg/ 24 hrs.